

## Prevalence of mamelons among chennai population - A crosssectional study

**Ashfaaq Ahmed**

*Undergraduate student*

*Saveetha Dental College and Hospitals*

*Saveetha Institute of medical and technical sciences*

*Saveetha university,*

*Chennai - 600077*

*Email ID: ashfaaq2002@gmail.com*

**Dr Reshma Poothakulath Krishnan**

*Senior Lecturer*

*Dept of Oral Pathology*

*Saveetha Dental College and Hospital*

*Saveetha Institute of medical and technical Sciences Saveetha*

*university, Chennai - 600077.*

[reshmapk.sdc@saveetha.com](mailto:reshmapk.sdc@saveetha.com)

**Dr. Sandhya Sundar**

*Senior Lecturer*

*Dept of Oral Pathology*

*Saveetha Dental College and Hospital*

*Saveetha Institute of medical and technical Sciences Saveetha*

*university, Chennai - 600077.*

[sandhyas.sdc@saveetha.com](mailto:sandhyas.sdc@saveetha.com)

---

### Abstract

**Background:** Mamelons are small projections on the incisal edge of the permanent incisor teeth. A newly erupted incisor shows protuberances on the incisal edge, separated by grooves which are known as mamelons. Mamelons are a three rounded bump on the edge of the tooth, it's made of enamel. Seen mostly on the incisal edge of the newly emerged permanent incisor teeth. In this study we tried to correlate the presence and absence of mamelons among different genders in the Chennai population. **Aim:** To evaluate the prevalence of mamelons on the incisor teeth among adolescents of chennai population **Materials and methods:** This study has been carried out among the undergraduate students of a private dental college in Chennai. The participants were between the age group 18-25 years. They were visually examined for the presence of mamelons. The data collected were tabulated and analyzed using Pearson's chi-square test using IBM SPSS software (version 23). **Results:** 24% of the study participants (n=12) showed the presence of mamelons which consisted of 8% females(n=4) and 16% males(n=8) in the age group 18-21 years. Results showed significantly higher prevalence in males when compared to females (chi-square, p value= 0.321). **Conclusion:** According to the study, more males have mamelons than females in the Chennai population. It is seen to persist more in the maxillary incisor teeth.

**Keywords:** Mamelons, Protuberance, Forensic evaluation, innovative technology, novel method

## **Introduction**

Mamelons is a three rounded protuberance on the edge of the incisal tooth, separated by grooves known as mamelon (1). It is made up of enamel, together they create a scalloped, wavy edge (1,2). It is easily noticeable on the permanent teeth of children, although it is possible for adults to have them. It erupts newly through the gums(3). It exists to help teeth break out through the gums. However it is agreed that they don't have any clinical significance.

They get smoothed out when the upper and lower front teeth come into contact, it may not go away if teeth are misaligned(4). It usually occurs when you have an open bite, in which the back and front teeth don't overlap each other vertically, hence the mamelons remain in adulthood(4,5). Mamelons are present on the maxillary central incisor, they are different, middle one is smallest and distal has a low shoulder, most mesial one has a raised shoulder.(4–6). Mamelons are helpful to differentiate primary and permanent dentition, since they are absent on the primary dentition. They are present in diseases like KBG syndrome(7). They do not have a dentin underneath.


Mamelons can be removed by shaving the teeth's edges, this is a form of cosmetic dentistry treatment, it is painless and does not require a local anesthetic because it does not contain any nerves, it is a quick and cheap procedure. It is removed for appearance reasons only, it does not cause any harm(6). Our team has extensive knowledge and research experience that has translate into high quality publications (8), (9), (10), (11), (12), (13), (14), (15), (16), (17), (18), (19), (20), (21), (22), (23), (24), (25), (26), (27). Thus this study is conducted to analyze the persistence of mamelons among different genders in the Chennai population.

## **Materials and methods**

This study has been carried out on 50 South Indian students in Saveetha dental college, Chennai to evaluate the incidence of mamelons for forensic evaluation. This study compares the prevalence of mamelons with relation to sex of same age group, for the age group of 18-25 years. The consent of the participants in this study was taken. Presence of the mamelons were observed through visual examination with the help of mouth mirror, the mamelons were present on both sides for some cases. The Chi-square tests were done using IBM SPSS software(version 23).

## **Results:**

A total of 50 participants were recruited in the study. After they were subjected to physical examination in the dental chair, it was found that 12 (24%) of them had shown the presence of mamelons. Males showed more presence of mamelons more frequently - 67% (8 nos.) compared to that of females - 33% (4 nos.) (p value= 0.321). The association between the presence of mamelons in different genders are graphically represented below as a bar chart in graph 1.



**Graph 1:** The bar graph represents the association between gender and presence of mamelons. X axis represents the gender and Y axis represents the number of participants. Blue denotes absence of mamelon and green denotes presence of mamelon. 16% of males showed the presence of mamelons whereas 34% of them showed absence of it. In females, 8% of the population showed the presence of mamelon and 42% did not have the mamelon. (Chi-square test; P value = .321, hence not significant)

**Discussion**

Mamelons are three rounded bumps or scallops formed on one of the three facial development lobes on the incisal edges of newly erupted incisors, separated by grooves(28). They help the teeth to cut through the gums more easily. It slowly wears off and is seen in adults in rare cases because of factors like delayed tooth eruption or occlusal inconsistency where the grind does not naturally grind, hence the mamelons remain in adulthood in that case. Mamelons are only enamel extensions, with no dentin underneath which pronounce them as translucent and more noticeable components(29). Mamelons are absent on the primary dentition, so they can be considered as important structures to differentiate the nature of dentition.

It is not evident on adult dentition in most cases, no cases of mamelons were seen in the age group above 50. This fact goes so well that mamelons are worn off with wear and tear.(30) It is seen better on the central incisor as compared to the lateral incisor, when the maxillary and mandibular teeth have an anterior open- bite relationship. The mamelons are present in adults where there is no functional contact between the maxillary and mandibular teeth. These are harmless and do not interfere with chewing

however it can be corrected for aesthetic purposes through treatments like teeth recontouring, cosmetic contouring and teeth reshaping(31).

In the present study also out of 50 only 12 individuals had the presence of mamelons where males had significantly higher presence than females(p value= 0.321). In a different study it is analyzed that persistence of mamelons were present more in females than males(32) which was different from the findings of the current study which shows that the males have more persistence to mamelons significantly than females. Drawback of this study is that the sample size was less and only one age group but we would like to carry further research on mamelons testing on a large scale of people with a vast age group.

### **Conclusion**

According to the study, males have a higher rate of presence of mamelons than females in the Chennai population. A standard protocol regarding training as well as conservative correction of mamelons for aesthetic purposes should be formulated in all dental institutions. Further studies should be done with a larger sample size to generalize the results.

### **ACKNOWLEDGEMENT:**

We thank Saveetha Dental College and Hospitals for providing us the support to conduct the study.

### **CONFLICT OF INTEREST:**

The author declares that there was no conflict of interest in the present study

### **SOURCE OF FUNDING:**

The present study was supported by the following agencies.

- Saveetha Dental College
- Saveetha Institute of Medical and Technical Sciences (SIMATS)
- Saveetha University
- Indo Vietnam Association, Pondicherry

### **References**

1. Alalawi M, Al-Jhany N. Frequency of Mamelons in Relation to Age, Gender and Occlusion among the Saudi Population [Internet]. Vol. 2, Arab Journal of Forensic Sciences & Forensic Medicine. 2020. p. 106–10. Available from: <http://dx.doi.org/10.26735/vglp5383>
2. Murray WHH. The Doom of Mamelons: A Legend of the Saguenay. 1888. 134 p.
3. Chassain K, Vrignaud A, Cesbron E, Bara-Passot C, Maillard H. Eczéma des mamelons responsable d'une galactocèle par hyperprolactinémie mécanique [Internet]. Vol. 146, Annales de Dermatologie et de Vénérologie. 2019. p. A210–1. Available from: <http://dx.doi.org/10.1016/j.annder.2019.09.314>

4. Inc. KN, Kernel Networks Inc. Antimicrobial Photodynamic Therapy on Teeth With Molar Incisor Hypomineralization [Internet]. Case Medical Research. 2019. Available from: <http://dx.doi.org/10.31525/ct1-nct03904641>
5. Rawlinson J, Carmalt JL. Extraction techniques for equine incisor and canine teeth [Internet]. Vol. 26, Equine Veterinary Education. 2014. p. 657–71. Available from: <http://dx.doi.org/10.1111/eve.12252>
6. Alamankany A. Riga-Fede disease associated with Fabry's disease and Niemann-Pick C disease in a boy with microcephaly: A case report. *Int J Health Sci* . 2021 Jan;15(1):56–8.
7. Morghen I, Ferri E. The KBG syndrome: Case report [Internet]. Vol. 1, Cases Journal. 2008. Available from: <http://dx.doi.org/10.1186/1757-1626-1-186>
8. Princeton B, Santhakumar P, Prathap L. Awareness on Preventive Measures taken by Health Care Professionals Attending COVID-19 Patients among Dental Students. *Eur J Dent*. 2020 Dec;14(S 01):S105–9.
9. Mathew MG, Samuel SR, Soni AJ, Roopa KB. Evaluation of adhesion of *Streptococcus mutans*, plaque accumulation on zirconia and stainless steel crowns, and surrounding gingival inflammation in primary molars: randomized controlled trial. *Clin Oral Investig*. 2020 Sep;24(9):3275–80.
10. Sridharan G, Ramani P, Patankar S, Vijayaraghavan R. Evaluation of salivary metabolomics in oral leukoplakia and oral squamous cell carcinoma. *J Oral Pathol Med*. 2019 Apr;48(4):299–306.
11. R H, Hannah R, Ramani P, Ramanathan A, Jancy MR, Gheena S, et al. CYP2 C9 polymorphism among patients with oral squamous cell carcinoma and its role in altering the metabolism of benzo[a]pyrene [Internet]. Vol. 130, Oral Surgery, Oral Medicine, Oral Pathology and Oral Radiology. 2020. p. 306–12. Available from: <http://dx.doi.org/10.1016/j.oooo.2020.06.021>
12. Antony JVM, Ramani P, Ramasubramanian A, Sukumaran G. Particle size penetration rate and effects of smoke and smokeless tobacco products - An invitro analysis. *Heliyon*. 2021 Mar 1;7(3):e06455.
13. Sarode SC, Gondivkar S, Sarode GS, Gadbail A, Yuwanati M. Hybrid oral potentially malignant disorder: A neglected fact in oral submucous fibrosis. *Oral Oncol*. 2021 Jun 16;105390.
14. Hannah R, Ramani P, WM Tilakaratne, Sukumaran G, Ramasubramanian A, Krishnan RP. Author response for “Critical appraisal of different triggering pathways for the pathobiology of pemphigus vulgaris—A review” [Internet]. Wiley; 2021. Available from: <https://publons.com/publon/47643844>
15. Chandrasekar R, Chandrasekhar S, Sundari KKS, Ravi P. Development and validation of a formula for objective assessment of cervical vertebral bone age. *Prog Orthod*. 2020 Oct 12;21(1):38.

16. Subramanyam D, Gurunathan D, Gaayathri R, Vishnu Priya V. Comparative evaluation of salivary malondialdehyde levels as a marker of lipid peroxidation in early childhood caries. *Eur J Dent*. 2018 Jan;12(1):67–70.
17. Jeevanandan G, Thomas E. Volumetric analysis of hand, reciprocating and rotary instrumentation techniques in primary molars using spiral computed tomography: An in vitro comparative study. *Eur J Dent*. 2018 Jan;12(1):21–6.
18. Ponnulakshmi R, Shyamaladevi B, Vijayalakshmi P, Selvaraj J. In silico and in vivo analysis to identify the antidiabetic activity of beta sitosterol in adipose tissue of high fat diet and sucrose induced type-2 diabetic experimental rats. *Toxicol Mech Methods*. 2019 May;29(4):276–90.
19. Sundaram R, Nandhakumar E, Haseena Banu H. Hesperidin, a citrus flavonoid ameliorates hyperglycemia by regulating key enzymes of carbohydrate metabolism in streptozotocin-induced diabetic rats. *Toxicol Mech Methods*. 2019 Nov;29(9):644–53.
20. Alsawalha M, Rao CV, Al-Subaie AM, Haque SKM, Veeraraghavan VP, Surapaneni KM. Novel mathematical modelling of Saudi Arabian natural diatomite clay. *Mater Res Express*. 2019 Sep 4;6(10):105531.
21. Yu J, Li M, Zhan D, Shi C, Fang L, Ban C, et al. Inhibitory effects of triterpenoid betulin on inflammatory mediators inducible nitric oxide synthase, cyclooxygenase-2, tumor necrosis factoralpha, interleukin-6, and proliferating cell nuclear antigen in 1, 2-dimethylhydrazine-induced rat colon carcinogenesis. *Pharmacogn Mag*. 2020;16(72):836.
22. Shree KH, Hema Shree K, Ramani P, Herald Sherlin, Sukumaran G, Jeyaraj G, et al. Saliva as a Diagnostic Tool in Oral Squamous Cell Carcinoma – a Systematic Review with Meta Analysis [Internet]. Vol. 25, *Pathology & Oncology Research*. 2019. p. 447–53. Available from: <http://dx.doi.org/10.1007/s12253-019-00588-2>
23. Zafar A, Sherlin HJ, Jayaraj G, Ramani P, Don KR, Santhanam A. Diagnostic utility of touch imprint cytology for intraoperative assessment of surgical margins and sentinel lymph nodes in oral squamous cell carcinoma patients using four different cytological stains. *Diagn Cytopathol*. 2020 Feb;48(2):101–10.
24. Karunagaran M, Murali P, Palaniappan V, Sivapathasundharam B. Expression and distribution pattern of podoplanin in oral submucous fibrosis with varying degrees of dysplasia – an immunohistochemical study [Internet]. Vol. 42, *Journal of Histotechnology*. 2019. p. 80–6. Available from: <http://dx.doi.org/10.1080/01478885.2019.1594543>
25. Sarode SC, Gondivkar S, Gadbaile A, Sarode GS, Yuwanati M. Oral submucous fibrosis and heterogeneity in outcome measures: a critical viewpoint. *Future Oncol*. 2021 Jun;17(17):2123–6.
26. Raj Preeth D, Saravanan S, Shairam M, Selvakumar N, Selestina Raja I, Dhanasekaran A, et al.

- Bioactive Zinc(II) complex incorporated PCL/gelatin electrospun nanofiber enhanced bone tissue regeneration. *Eur J Pharm Sci*. 2021 May 1;160:105768.
27. Prithviraj N, Yang GE, Thangavelu L, Yan J. Anticancer Compounds From Starfish Regenerating Tissues and Their Antioxidant Properties on Human Oral Epidermoid Carcinoma KB Cells. In: PANCREAS. LIPPINCOTT WILLIAMS & WILKINS TWO COMMERCE SQ, 2001 MARKET ST, PHILADELPHIA ...; 2020. p. 155–6.
  28. Praiss IL, Tannenbaum KA, Gelder-Kogan CA. Quality of dental care--the role of third-party payers: a literature review. *Med Care Rev*. 1978 Dec;35(11):1211–33.
  29. Kent Obermann W. Evaluation of Mamelons: Their Incidence and Frequency Distributions with Regard to Race and Sex. 1982. 150 p.
  30. Abdel-Gaber R. *Syphacia obvelata* (Nematode, Oxyuridae) infecting laboratory mice *Mus musculus* (Rodentia, Muridae): phylogeny and host-parasite relationship. *Parasitol Res*. 2016 Mar;115(3):975–85.
  31. Singh G. Cosmetic Contouring in Orthodontics [Internet]. *Textbook of Orthodontics*. 2007. p. 313–313. Available from: [http://dx.doi.org/10.5005/jp/books/10936\\_28](http://dx.doi.org/10.5005/jp/books/10936_28)
  32. Rauber GB, Bernardon JK, Vieira LCC, Baratieri LN. Evaluation of a technique for color correction in restoring anterior teeth. *J Esthet Restor Dent*. 2017 Sep;29(5):309–16.