

## Efficiency Of *Pseudomonas Aeruginosa* In Bioremediation Of Chlorpyrifos Toxicity

Hassanin Sabah Hashim <sup>1</sup>, Ameera O. Hussain Al-Janabi <sup>2</sup> and jassim abdulaziz humadi alfalahy

<sup>1</sup>Department Biotechnology, College of Biotechnology, University of Al-Qasim Green, AL-Qasim Babylon, IRAQ.

<sup>2</sup>Department of Environmental, College of Environmental Sciences, University of Al-Qasim Green, Qasim Babylon, IRAQ.

---

### Abstract:

The mode of action involves inhibiting acetyl-cholinesterase leading to the accumulation of acetylcholine causing neurotoxicity. It is being transported by circulation far away from the site of application leading to pollution of the environment. Study aimed at investigating the efficiency of *Pseudomonas aeruginosa* in bioremediation of chlorpyrifos from contaminated soil in Iraq. HPLC technology was used to estimate the percentage of removal achieved by bacteria, and it is considered one of the advanced modern techniques in measuring ingredient active in solutions. In conclusion from the present study the results of the HPLC analysis showed that *Pseudomonas aeruginosa* is highly efficient in the analysis of chlorpyrifos.

**Keywords:** Bioremediation, organophosphate, Chlorpyrifos, *Pseudomonas aeruginosa*, HPLC.

---

### Introduction

Chlorpyrifos (O,O-diethyl-O-3,5,6-trichlor-2-pyridyl phosphorothioate; CPF) is a broad spectrum organophosphate insecticide (OP) that is commercially used to control insects especially termite (Rusyniak and Nañagas, 2004). It was first developed by the Germans in the 1930s and first introduced into the marketplace in 1965, has been widely used globally as an insecticide to control crop pests in agriculture, reduce household pests such as termites, reduce insect damage to turf on lawns, and for mosquito control (Chishti, et al., 2013; Eaton, et al., 2008).

A variety of studies have shown that exposure to CPF elicits acute toxicity through inhibition of acetylcholinesterase (AChE) throughout the central and peripheral nervous systems, leading to acetylcholine accumulation and long-term stimulation of cholinergic receptors (Singh et al., 2009). Chlorpyrifos works basically the same way both in insects and other animals including humans through impairment of nervous system. During development can cause persisting neurobehavioral dysfunction, even with low doses that do not elicit acute toxicity (Koly and Khan, 2006).

The fate of chlorpyrifos is affected not only by its own physicochemical properties, but also by characteristics of the soil, management practices and environmental conditions (Muhamad, et al., 2010) Its

persistence increases with decreased temperature, decreased pH, and decreased light. The half-life of chlorpyrifos showed that it remained stable even after 12 months in the soil (Koly and Khan, 2019) and Chlorpyrifos is moderately soluble in water, the half-life of chlorpyrifos in water ranges from 35 to 78 days (Chishti, et al., 2013).

have short to moderate persistence in the environment as a result of several non-biological methods including volatilization, photolysis, abiotic hydrolysis, and also by microbial degradation that might occur concurrently (Koly and Khan, 2019). Chlorpyrifos has been of a great concern due to persistence, toxicity and accumulation in soils and ground waters (Abdul-jabbar., et al 2017) and It was observed may be causes altered the structure of the gills and liver of fishes (Deb and Das, 2013).

The unsafe misuse of pesticides has caused widespread harm to human health and the environment, because it is one of the most common causes of poisoning around the world (Lu et al, 1999) as well as the appearance of its residues in crops and soil, and has proven many Reports, including the Environmental Protection Organization, that pesticides cause many types of cancer (Helfrich, et al., 2009) When pesticides are dispersed in the environment, they become pollutants, with ecological effects that require remediation (Ortiz-Hernández and Sánchez-Salinas, 2011).

## **2-Materials and methods**

### **2-1 Sample collection**

The soil samples used for the isolation of pesticide degrading bacteria were collected from agricultural fields, residential buildings, and Garden yards from the Iraq-Al Diwaniyah where chlorpyrifos pesticide is used extensively. a total of 50 samples were collected at a depth of 5cm to 10cm from different regions. Commercial chlorpyrifos (48%) pesticide was procured from local pesticide shop . Stock solution was prepared and stored in refrigerator for further use. The samples were collected using sterile petri dish and plain tube and 5 grams of soil samples were taken and added Normal line to it, for the purpose of purification and isolation of bacteria.

### **2-3 Preparation of stock pesticide**

To estimate the amount of residual chlorpyrifos concentrations after removal by bacterial isolates. Standard was prepared for the active ingredient chlorpyrifos by selecting more than one manufacturer of the pesticide containing the active ingredient chlorpyrifos at a concentration of 48%, then 5 mg was taken from each package and analyzed by high performance chromatography according to (Mauldin et al., 2006)

### **2-4 Isolation and identification of bacteria**

Prepare the soil suspensions by taking 1 gram of soil, and dissolve in 10 ml of normal saline, then mix by the vortex for 10 minutes and centrifuge for 5 min at, and then filter with a 0.4 volume filter and use pseudomonas agar medium from (Himedia Company, India) to isolate the bacteria that analysis the pesticides. The culture medium consisting of Gelatin peptone 16.00 g / L, Tryptone 10,000 g / L, potassium sulphate 10.00 g / l, magnesium chloride 1.4 g / l, agar 11.00 g / l, and sterilized by autoclaving at 15lbs (121° C) for 15 min, cool to 45-50 ° C and aseptically add sterile rehydrated contents of one vial of either celrinix supplement (FD029) or CFC supplement (FD036) as desired. The pH was set to 7.1 at 25 ° C. mix well and pour into sterile petri dish plates for selective isolation of pseudomonas species.

The cultivated bacterial isolates were incubated on the nutrient medium for 24 hours at a temperature of 37 ° C and the bacteria were diagnosed by a modern device called VITEK2 SYSTEM from the company (Biomérieux, France). VITEK2 SYSTEM was examined against three bacteria as shown in (table. 1).

### 2-5 Treating the medium with the pesticide

Nutrient broth is prepared for the purpose of treatment with chlorpyrifos and consists of peptone 5 g / liter, sodium chloride 5 g / liter, meat extract B 1.5 g / liter, yeast extract 1.5 g / liter. Addition 800 ml of distilled water and dissolve 10.2 grams from the medium, then heat the mixture by microwave and sterilize with an autoclave at 15lbs (121° C) for 15min and then cool to 45° C.

### 2-6 Biodegradation of chlorpyrifos

add 3 ml of water and use a 0.4 volume filter to sterilize the water with a syringe, add 200µl of chlorpyrifos, and mix well and add the last mixture to the liquid medium in order to inoculate the medium with the isolated and diagnosed bacteria . The residual concentrations were assessed and measured after removal with Hplc and uninoculated flasks served as controls. Biodegradation experiments were carried out in 250 mL Erlenmeyer flasks containing 60 mL of medium with 200 µL of CP. The mixture were pour in cup urin 50 ml and inoculated with pseudomonas aeruginosa and incubated at 37°C. Samples were taken at periodic intervals and the residual pesticide concentration was determined using HPLC. Un-inoculated cup served as controls.


### 3-Result and discussion

pseudomonas spp. is highly efficient in the biological removal of organic phosphorous pesticides (chlorpyrifos) at a concentration of 120 µg/l, thus the possibility of using it in the bioremediation of pollutants. P.aeruginosa strain removal 99% of CP with at 6 days. there were probability of diagnostic bacterial isolates by Vitek 2 Compact system(P.aeruginosa 99%,P. putida 99%,P. fluorescence 94%), the residual chlorpyrifos concentration was 1.10 to 0.00 µg /mL ., (table.2) while P.putida (fig.3) and P.fluorescence (fig.4) degraded CP within 6 days from 0.08µg/mL to 0.002µg/mL and 0.26µg/mL to 0.118µg/mL. The P.aeruginosa could rapidly degrade chlorpyrifos compare with other species during the logarithmic phase of growth. The percentage degradation in first 6 day was higher and rapidly increased thereafter to 100% within 12 day compare with control (Fig. 2). These results are consistent with many studies and research that have confirmed the ability of microorganisms (bacteria and fungi) to consume a wide range of pesticides and in most cases their ability to consume one or more compounds as a source of carbon and energy ( Ortiz-Hernández and Sánchez-Salinas,2010; Chishti, et al., 2013) The strain could utilize CP as the sole source of carbon for growth. In HPLC analysis of sample uninoculated (without bacteria) with 200 µl CP, two peaks appeared in HPLC separation (Fig.1), which were further identified as CP by comparing with their authentic standard. HPLC analysis showed that CP could be degraded by P. aeruginosa to TCP and diethylthiophosphoric acid.


**Table 2. the percentages of residual concentrations after treatment with for chlorpyrifos**

Bacterial sample	Concentrate Of chlorpyrifos	Residual Concentrate After 6 days	Residual Concentrate after 12 days	Control

<b>P.aeruginosa</b>	120 µg/ml	1.10 µg/ml	0.000 µg/ml	63.40 µg/ml
<b>p.putida??</b>		0.08 µg/ml	0.002 µg/ml	
<b>p.fluorescence</b>		0.26 µg/ml	0.118 µg/ml	



**Figure (1):** absorbance and spectrum of chlorpyrifos and the control (sample uninoculated) treated with the pesticide without bacteria.



**Figure (2):** growth kinetic for *P. aeruginosa* on broth media with chlorpyrifos as carbon source and energy.




Figure (3): I peaks Chlorpyrifos (CP) degradation by *Pseudomonas putida* strain. 1)




Figure (4): Illustrates peaks Chlorpyrifos (CP) degradation by *Pseudomonas fluorescence*

Only a few organisms capable of degrading chlorpyrifos have been reported till date. The maximum rate of chlorpyrifos removal was observed from 6 to 12 days, which coincided with the period of maximum proliferation. This might indicate that the *pseudomonas aeruginosa* utilized CP as a source of energy during its growth phase. It was noted that on reaching lower concentrations of residual pesticide in the medium, the degradation rate decreased significantly as previously observed by.

The minimum incubation time required for complete removal of chlorpyrifos was 6 h by *Cupriavidus* sp. DT-1 in a liquid medium (Lu et al., 2013). Most of the common CP degrading bacteria were reported to degrade the pesticide within 48 h in liquid media (Gómez et al., 2007). Chlorpyrifos has been shown to be degraded cometabolically in liquid media by bacteria (Jaiswal, et al., 2017) *Pseudomonas aeruginosa* is the most common Gram negative bacterium found in soil. Isolates of this bacterium have been found to have potential to degrade

chlorpyrifos (Fulekar and Geetha, 2008). Since *Pseudomonas aeruginosa* capable of degradation of various OPs have great potential for bioaugmentation of soil (Cycoń et al, 2017); these strain of bacteria with its high degradation potential, fast growth, minimal nutrient requirements and tolerance to high concentrations of pesticide can play a vital role in decontamination of polluted soil. The rapid removal of CP from liquid medium within 6 day by the *p.aeruginosa* strain thus could be one of the fastest reported for effective CP degradation and first of its kind amongst the *Pseudomonads*. *P.aeruginosa* is a known plant growth promoter (Trinh et al., 2018), and this combined with the reduced time for decontamination of CP makes it a promising candidate for pesticide removal from agricultural lands and also for treating pesticide industry effluents.

To demonstrate the capacity of the recombinant *P. aeruginosa* strain to degrade CP, the degradation experiments were performed by inoculating the recombinant strain into a nutrient broth medium supplemented with CP. High-performance liquid chromatography (HPLC) analysis indicated that CP were completely degraded within 6 and 12 days respectively. Moreover, the concentration of 3,5,6- trichloro-2-pyridinol (TCP) in the medium increased gradually with the decrease in CP concentration. CP was degraded quickly in the first 6 days, which accounted for 48% of the amount of the initially added pesticides respectively. Maybe a reduction in the degradation rate after 12 days may be due to the accumulation of TCP, which have antimicrobial activity and are toxic to the bacterial growth and metabolism (Singh et al., 2004 ; Nguyen et al., 2014)

#### 4. CONCLUSION

The present study shows that use of the *pseudomonas aeruginosa* culture could effectively degrade chlorpyrifos pesticide. Removal of xenobiotic compounds which are a major threat to environmental pollution are of major concern as they enter the food chain and are the major causes of various diseases. The study shows that the bacterial isolate is able to use chlorpyrifos as sole source of carbon. Therefore these bacteria culture can be used effectively for Bioremediation of contaminated sites. Future studies aim in studying the biotechnology of pesticide degradation and its significance in field conditions.

#### Reference

1. Rusyniak, D.E. and K.A. Nañagas. Organophosphate poisoning. in Seminars in neurology. 2004. Copyright© 2004 by Thieme Medical Publishers, Inc., 333 Seventh Avenue, New ...
2. Chishti, Z., et al., Microbial degradation of chlorpyrifos in liquid media and soil. *J Environ Manage*, 2013. **114**: p. 372-80.
3. Eaton, D.L., et al., Review of the toxicology of chlorpyrifos with an emphasis on human exposure and neurodevelopment. *Critical reviews in toxicology*, 2008. **38**(sup2): p. 1-125.
4. Singh, P., et al., Biosurfactant production by *Pseudomonas* sp. and its role in aqueous phase partitioning and biodegradation of chlorpyrifos. *Letters in applied microbiology*, 2009. **49**(3): p. 378-383.
5. Koly, F.A. and R.A. Khan,. Biodegradation of organophosphorous pesticide: chlorpyrifos. *Scientific Review*, 2019. **5**(1): p. 8-18.
6. Muhamad, H., et al., Downward movement of chlorpyrifos in the soil of an oil palm plantation in Sepang, Selangor, Malaysia. *Journal of Oil Palm Research*, 2010. **22**(1): p. 721-728.
7. Abdul-jabbar Abbas Ali., et al.,. Treatment of Chlorpyrifos Pesticide in Water Using Nano Scales Zero Valent Iron (nZVI). *Iraqi Journal of Science and Technology*, 2017. **8**(1): p. 91-97.
8. Deb, N. and S. Das, Chlorpyrifos toxicity in fish: a review. *Current World Environment*, 2013. **8**(1): p. 77.

9. Lu, Y.-T., N.G. Love, and C.L. Grady Jr, Microscopic methods for distinguishing among three cell types in TOL plasmid-carrying *Pseudomonas putida* cultures. *FEMS microbiology letters*, 1999. **173**(1): p. 195-201.
10. Helfrich, L.A., et al., *Pesticides and aquatic animals: a guide to reducing impacts on aquatic systems*. 2009.
11. Ortiz-Hernández, M.L. and E. Sánchez-Salinas, Biodegradation of the organophosphate pesticide tetrachlorvinphos by bacteria isolated from agricultural soils in México. *Revista internacional de contaminación ambiental*, 2010. **26**(1): p. 27-38.
12. Nyende, A., et al. BIODEGRADATION OF DIAZINON AND METHOMYL PESTICIDES BY WHITE ROT FUNGI FROM SELECTED HORTICULTURAL FARMS IN RIFT VALLEY AND CENTRAL PROVINCES, KENYA. in *JKUAT ANNUAL SCIENTIFIC CONFERENCE PROCEEDINGS*. 2010.
13. de Sousa Fragoeiro, S.I., *Use of fungi in bioremediation of pesticides*. 2005, Cranfield University at Silsoe.
14. Jaiswal, S., et al., Bioremediation of Chlorpyrifos Contaminated Soil by Microorganism. *International Journal of Environment, Agriculture and Biotechnology*, 2017. **2**(4): p. 238833.
15. Mauldin, R.E., et al., A simple HPLC method for the determination of chlorpyrifos in black oil sunflower seeds. *Journal of liquid chromatography & related technologies*, 2006. **29**(3): p. 339-348.
16. Lu, P., et al., Biodegradation of chlorpyrifos and 3, 5, 6-trichloro-2-pyridinol by *Cupriavidus* sp. DT-1. *Bioresource technology*, 2013. **127**: p. 337-342.
17. Gómez, M.J., et al., The environmental fate of organic pollutants through the global microbial metabolism. *Molecular Systems Biology*, 2007. **3**(1): p. 114.
18. Fulekar, M. and M. Geetha, Bioremediation of Chlorpyrifos by *Pseudomonas aeruginosa* using scale up technique. *J Appl Biosci*, 2008. **12**: p. 657-60.
19. Cycoń, M., A. Mroziak, and Z. Piotrowska-Seget, Bioaugmentation as a strategy for the remediation of pesticide-polluted soil: A review. *Chemosphere*, 2017. **172**: p. 52-71.
20. Trinh, C.S., et al., Evaluation of the plant growth-promoting activity of *Pseudomonas nitroreducens* in *Arabidopsis thaliana* and *Lactuca sativa*. *Plant cell reports*, 2018. **37**(6): p. 873-885.
21. Singh, B.K., et al., Biodegradation of chlorpyrifos by *Enterobacter* strain B-14 and its use in bioremediation of contaminated soils. *Appl. Environ. Microbiol.*, 2004. **70**(8): p. 4855-4863.
22. Nguyen, T.P.O., et al., Genetic and metabolic analysis of the carbofuran catabolic pathway in *Novosphingobium* sp. KN65. 2. *Applied microbiology and biotechnology*, 2014. **98**(19): p. 8235-8252.